-

Junior-Senior Individual Test

Directions: Please answer all questions on the answer sheet provided. All answers must be written legibly in the correct blanks on the answer sheet and in simplest form. **Exact** answers are to be given unless otherwise specified in the question. No units of measurement are required. Each problem has the same point-value.

- 1. Find the sum of all distinct values of x such that $\left[\log_k(x^2)\right]\left(\log_{12}k\right) = 2$.
- 2. Let $i = \sqrt{-1}$. Then $-2i^2 + (\sqrt{-4})(\sqrt{4}) (\sqrt{-3})(\sqrt{-3}) 2i^5 = a + hi$, where a and h are real numbers. Find

the value of (3a+2b).

- 3. If x is an integer, find the sum of all distinct values of x such that $\frac{x-4}{x-9} 3 \ge 0$.
- $\underline{4}$, In the diagram. \underline{A} \underline{B} and \underline{D} lie on the circle with center \underline{O}



- 10. Find the value of $\log_{27} \left(9 \left(\frac{1}{27} \right)^{-2} \right)$. Give your answer as a fully reduced **improper** fraction.
- 11. Find the eighth term of an arithmetic progression whose first term is 3 and whose 31st term is 73. Give your answer as a fully reduced **improper** fraction.
- 12. Suppose that $\frac{8!}{3!k!}$ = 56. Find the value of k.
- 13 When 1 2 3 1 and 5 are substituted for x in a notinomial expression for x the results are respectively

Name:	Team Code:	
2014 John O'Bryan Mathematical Competition Junior/Senior Individual Test		
Note All anguer	against ha sivittan lagibly in the compat blanks on the charge sheet and in him less feed	
1		
<u>a</u>		
· Y ————————————————————————————————————		
• * •		
•		
Exact ansv are requir	vers are to be given unless otherwise specified in the question. No units of measurement ed. Each problem has the same point-value.	
1.	, 11	

.

Name:	ANSWERS	Team Code:
·	2014 John O'Bryan M	athematical Competition
•		
7		
^		
E) at :	
<u> </u>	•	
	•	
(
-	₹- <u>-</u> -	